arXiv:1701.01154v1 [cs.IT] 8 Dec 2016

Perfect Sequences and Arrays over the Unit Quaternions[1]

Sam Blake
School of Mathematical Sciences, Monash University, Australia

We introduce several new constructions for perfect periodic autocorrelation sequences and arrays over the unit quaternions. This paper uses both mathematical proofs and computer experiments to prove the (bounded) array constructions have perfect periodic autocorrelation. Furthermore, the first sequence construction generates odd-perfect sequences of unbounded lengths, with good ZCZ.

The quaternions were discovered by the Irish mathematician Sir William Rowan Hamilton in 1843. Hamilton was interested in the connection between complex numbers and 2-dimensional geometry. He tried in vain to extend the complex numbers to $\mathbb{R}^3$, only years later would it be discovered that there is no 3-dimensional normed division algebra. Hamilton's breakthrough came when he extended the complex numbers to 4 dimensions. The quaternions are a 4-dimensional non-commutative division algebra. They can be thought of as an extension of the complex numbers, where

$$\boldsymbol{i}^2 = \boldsymbol{j}^2 = \boldsymbol{k}^2 = \boldsymbol{ijk} = -1,$$

and they multiply as follows (row $\times$ column)

$\times$	1	$\boldsymbol{i}$	$\boldsymbol{j}$	$\boldsymbol{k}$
1	1	$\boldsymbol{i}$	$\boldsymbol{j}$	$\boldsymbol{k}$
$\boldsymbol{i}$	$\boldsymbol{i}$	-1	$\boldsymbol{k}$	$-\boldsymbol{j}$
$\boldsymbol{j}$	$\boldsymbol{j}$	$-\boldsymbol{k}$	-1	$\boldsymbol{i}$
$\boldsymbol{k}$	$\boldsymbol{k}$	$\boldsymbol{j}$	$-\boldsymbol{i}$	-1

Given the quaternions, $\mathbf{q}_1 = a + b\boldsymbol{i} + c\boldsymbol{j} + d\boldsymbol{k}$, where $a, b, c, d \in \mathbb{R}$, $\mathbf{q}_1 = a_1 + b_1\boldsymbol{i} + c_1\boldsymbol{j} + d_1\boldsymbol{k}$ where $a_1, b_1, c_1, d_1 \in \mathbb{R}$, and $\mathbf{q}_2 = a_2 + b_2\boldsymbol{i} + c_2\boldsymbol{j} + d_2\boldsymbol{k}$ where $a_2, b_2, c_2, d_2 \in \mathbb{R}$, we have

$$\begin{aligned}
\mathbf{q}_1 + \mathbf{q}_2 &= (a_1 + a_2) + (b_1 + b_2)\boldsymbol{i} + (c_1 + c_2)\boldsymbol{j} + (d_1 + d_2)\boldsymbol{k} \\
\mathbf{q}_1\mathbf{q}_2 &= (a_1a_2 - b_1b_2 - c_1c_2 - d_1d_2) + (a_2b_1 + a_1b_2 - c_2d_1 + c_1d_2)\boldsymbol{i} + \\
&\quad (a_2c_1 + a_1c_2 + b_2d_1 - b_1d_2)\boldsymbol{j} + (a_2d_1 + a_1d_2 - b_2c_1 + b_1c_2)\boldsymbol{k} \\
\mathbf{q}^* &= a - b\boldsymbol{i} - c\boldsymbol{j} - d\boldsymbol{k} \\
\|\mathbf{q}\| &= \sqrt{\mathbf{q}^*\mathbf{q}} = \sqrt{a^2 + b^2 + c^2 + d^2} \\
\mathbf{q}^{-1} &= \frac{\mathbf{q}^*}{\|\mathbf{q}\|^2}
\end{aligned}$$

Over the quaternions, the notation $\frac{\mathbf{q}_1}{\mathbf{q}_2}$ is ambiguous, it could be left division, $\mathbf{q}_2^{-1}\mathbf{q}_1$, or right division, $\mathbf{q}_1\mathbf{q}_2^{-1}$.

If $\|\mathbf{q}\| = 1$, then $\mathbf{q}$ is a *unit* quaternion. All the sequences and arrays defined in this chapter are restricted to *simple unit* quaternions, which are given by $-1, 1, -\boldsymbol{i}, \boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{j}, -\boldsymbol{k}, \boldsymbol{k}$.

Moxey et al.[14] observed that as multiplication over the quaternions is non-commutative, the definition of autocorrelation becomes ambiguous. One could define a *right* correlation: ${\theta^{\text{right}}}_{\mathbf{s}}(\tau) = \sum_{i=0}^{n-1} s_i s_{i+\tau}^*$, and a *left* correlation: ${\theta^{\text{left}}}_{\mathbf{s}}(\tau) = \sum_{i=0}^{n-1} s_{i+\tau}^* s_i$. Thus we have two correlation measures over the quaternions. These two correlations are not always equal.

Example 0.1. We compute the left and right autocorrelations for the sequence, $\mathbf{s} = [\boldsymbol{i}, -\boldsymbol{j}, -1, -1, \boldsymbol{k}, 1]$, and show that the left and right autocorrelations of $\mathbf{s}$ are not equal for all shifts, τ:

τ	${\theta^{\text{left}}}_{\mathbf{s}}(\tau)$	${\theta^{\text{right}}}_{\mathbf{s}}(\tau)$
0	6	6
1	$\boldsymbol{k} - \boldsymbol{i} + \boldsymbol{j} + 1$	$3\boldsymbol{k} - \boldsymbol{i} + \boldsymbol{j} + 1$
2	$\boldsymbol{k} - \boldsymbol{i} + 3\boldsymbol{j} - 1$	$\boldsymbol{k} - \boldsymbol{i} + \boldsymbol{j} - 1$
3	-2	-2
4	$-\boldsymbol{k} + \boldsymbol{i} - 3\boldsymbol{j} - 1$	$-\boldsymbol{k} + \boldsymbol{i} - \boldsymbol{j} - 1$
5	$-\boldsymbol{k} + \boldsymbol{i} - \boldsymbol{j} + 1$	$-3\boldsymbol{k} + \boldsymbol{i} - \boldsymbol{j} + 1$

A sequence is *left perfect* if ${\theta^{\text{left}}}_{\mathbf{s}}(\tau) = 0$ for all off-peak shifts. Similarly, for *right perfect* sequences.

Kuznetsov[7] showed that a sequence is left perfect if and only if the sequence is right perfect. Thus, a left or right perfect sequence is simply referred to as a *perfect sequence*.

Perfect quaternion sequences and arrays have a very short history. To the best of the authors knowledge, the first appearance in the literature of quaternion correlations was by Sangwine and Ell[16] in 1999, where the Fourier transform of colour images was computed via a quaternion Fourier transform. The autocorrelation was computed using the explicit (signal processing) definition, which is quadratic in complexity.

In 1999 Leukhin et al.[10] described an application of quaternion sequences to optical signal processing involving photon echoes. There, a quaternion description of the polarisation state of the optical excitation is natural, and so is the quantum interaction within the material being analysed. Leukhin et al. discovered the first perfect sequence over the quaternions:
$\Big[1 + \boldsymbol{j}, 1 + \boldsymbol{j}, 1 + \boldsymbol{j}, 1 + \boldsymbol{j}, -\frac{1}{2} + \frac{\sqrt{3}\boldsymbol{i}}{2} - \frac{\boldsymbol{j}}{2} + \frac{\sqrt{3}\boldsymbol{k}}{2}, -\frac{1}{2} - \frac{\sqrt{3}\boldsymbol{i}}{2} - \frac{\boldsymbol{j}}{2} - \frac{\sqrt{3}\boldsymbol{k}}{2},$
$1 + \boldsymbol{j}, -\frac{1}{2} - \frac{\sqrt{3}\boldsymbol{i}}{2} - \frac{\boldsymbol{j}}{2} - \frac{\sqrt{3}\boldsymbol{k}}{2}, -\frac{1}{2} + \frac{\sqrt{3}\boldsymbol{i}}{2} - \frac{\boldsymbol{j}}{2} + \frac{\sqrt{3}\boldsymbol{k}}{2}\Big]$

In 2001 Pei et al.[15] described an algorithm for the quaternion Fourier transform, convolution, and correlation by a 2D complex fast Fourier transform.

In 2010 Kuznetsov et al.[8] used the known method of taking the product of sequences of coprime lengths to construct a quaternion sequence of length $5 \times 7 \times 9 \times 11 \times 13 \times 16 \times 17 \times 19 \times 23$. The sequences of smaller (odd) lengths were found by computer search, and of the form $[1, \boldsymbol{j}, \mathbf{s}, \mathbf{s}_r, \boldsymbol{j}, 1, \mathbf{q}]$, where $\mathbf{s}$ is a simple quaternion sequence and $q = \frac{a+b\boldsymbol{i}+c\boldsymbol{j}+d\boldsymbol{k}}{2}$, where $a, b, c, d = \pm 1$. The even length sequence is a Frank sequence[5] of length 16 over 4 roots of unity. Based on the length of the product sequence, Kuznetsov et al. conjectured that these sequences exist for unbounded lengths. We believe this conjecture is questionable. If the product sequences exist for unbounded lengths, then the individual sequences which form the product sequence must exist for unbounded lengths. The longest sequence found by computer search is only of length 23 over an alphabet with 23 members, which far less than the square of the number of members in the sequence alphabet, 529 (see Mow's conjecture in Chapter 1). Stronger evidence for the conjecture could, for example, be the construction of sequences longer than 529 which is not the product of smaller perfect sequences.

In 2012 Acevedo et al.[1] discovered a link between Lee sequences[9] and perfect quaternion sequences of lengths up to 98 which were discovered by the author in 2009[17]. As the Lee sequences exist for unbounded lengths, the corresponding sequences discovered by the author and subsequently generalised by Acevedo et al. are of unbounded length. While these sequences are over the quaternions, the single appearance of $\boldsymbol{j}$ and no appearance of $-\boldsymbol{j}$, $-\boldsymbol{k}$, and $\boldsymbol{k}$ suggests that for long sequences they are best thought of as *near*-quaternion sequences, as the frequency of each member in the sequence alphabet asymptotically approaches a sequence over 4 roots of unity.

Following the paper of Acevedo et al.[1], Acevedo and Jolly[2] extended the method of Arasu and de Launey[3] for constructing perfect arrays of unbounded size over 4 roots of unity to quaternions.

In the following constructions we make frequent use of a well-known property of complex numbers.

Lemma 0.2. *Let $m \in \mathbb{N}$ and $c \neq 0 \mod 4$, then* $\sum_{n=0}^{4m-1} \boldsymbol{i}^{cn} = 0$.

Proof. As $\boldsymbol{i}^{cn} = \boldsymbol{i}^{cn \bmod 4}$, the terms in the summation have a period of 4. Thus

$$\sum_{n=0}^{4m-1} \boldsymbol{i}^{cn} = m \sum_{n=0}^{3} \boldsymbol{i}^{cn} = \begin{cases} m\left(\boldsymbol{i}^0 + \boldsymbol{i}^1 + \boldsymbol{i}^2 + \boldsymbol{i}^3\right) = m(1 + \boldsymbol{i} - 1 - \boldsymbol{i}) = 0 & \text{if} \quad c = 1 \bmod 4 \\ m\left(\boldsymbol{i}^0 + \boldsymbol{i}^2 + \boldsymbol{i}^0 + \boldsymbol{i}^2\right) = m(1 - 1 + 1 - 1) = 0 & \text{if} \quad c = 2 \bmod 4 \\ m\left(\boldsymbol{i}^0 + \boldsymbol{i}^3 + \boldsymbol{i}^2 + \boldsymbol{i}^1\right) = m(1 - \boldsymbol{i} - 1 + \boldsymbol{i}) = 0 & \text{if} \quad c = 3 \bmod 4 \end{cases}$$

∎

Of course, Lemma 0.2 holds if $\boldsymbol{i}$ is replaced with $\boldsymbol{j}$, or $\boldsymbol{k}$. This summation is the quaternion equivalent of the Gaussian sums for roots of unity for perfect sequences. That is, we reduce autocorrelation summations to terms containing this summation in order to show they are perfect.

1 Sequences of lengths 10, 14, 18, 26, 30, 38, 42, 50, 54, 62, 74, 82, 90, and 98

The work presented in this section predates the work of Acevedo et al.[1] (See [17]).

In 2008 we used the Monash Campus Cluster[12] to perform exhaustive searches for perfect periodic autocorrelation sequences over n-tuples of $[-\boldsymbol{i}, \boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{j}, -\boldsymbol{k}, \boldsymbol{k}]$. As cancellation occurs pairwise, perfect sequences over this alphabet only exist for even lengths. The exhaustive searches for small lengths produced many sequences, for example

length	example sequence
4	$[-\boldsymbol{k}, \boldsymbol{i}, -\boldsymbol{k}, -\boldsymbol{i}]$
6	$[\boldsymbol{j}, \boldsymbol{k}, -\boldsymbol{j}, \boldsymbol{k}, \boldsymbol{j}, -\boldsymbol{i}]$
8	$[\boldsymbol{k}, \boldsymbol{i}, \boldsymbol{k}, -\boldsymbol{j}, -\boldsymbol{k}, \boldsymbol{i}, -\boldsymbol{k}, -\boldsymbol{j}]$
10	$[-\boldsymbol{i}, \boldsymbol{k}, \boldsymbol{i}, -\boldsymbol{k}, \boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{i}, -\boldsymbol{k}, \boldsymbol{i}, \boldsymbol{k}]$
12	–
14	$[-\boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{i}, \boldsymbol{j}, \boldsymbol{i}, \boldsymbol{j}, \boldsymbol{k}, \boldsymbol{j}, \boldsymbol{i}, \boldsymbol{j}, \boldsymbol{i}, -\boldsymbol{j}, -\boldsymbol{i}, \boldsymbol{j}]$
16	$[-\boldsymbol{i}, -\boldsymbol{i}, -\boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{i}, \boldsymbol{k}, -\boldsymbol{k}, -\boldsymbol{j}, -\boldsymbol{i}, \boldsymbol{i}, \boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{i}, -\boldsymbol{k}, \boldsymbol{k}, -\boldsymbol{j}]$

After looking through hundreds of small length sequences a symmetry was observed; many of the perfect sequences have the structure

$$\mathbf{q} = [-\boldsymbol{i}, \mathbf{s}, \boldsymbol{k}, \mathbf{s}_r]\,,$$

where $\mathbf{s} = [\alpha_0, \alpha_1, \cdots, \alpha_{n-1}] \times [\boldsymbol{j}, \boldsymbol{i}, \boldsymbol{j}, \boldsymbol{i}, \cdots, \boldsymbol{j}, \boldsymbol{i}]$, $\alpha_i = +1$ or -1, n is odd, and $\mathbf{s}_r$ is the reverse of $\mathbf{s}$. We refer to $\mathbf{s}$ as the sub-sequence of $\mathbf{q}$. Using this symmetry we can cut the search space for a sequence of length n from 6^n to $2^{(n-2)/2}$, and thus we can find significantly

longer sequences.

14	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{k},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
18	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
26	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
30	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},$ $\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
38	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},$ $-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
42	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{k},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},$ $\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
50	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},\boldsymbol{j},$ $-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
54	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},$ $\boldsymbol{k},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
62	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},$ $\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},$ $\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
74	$[-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},$ $\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},$ $-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j}]$
82	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},$ $-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{k},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},$ $-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},$ $-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
90	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},$ $-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},$ $-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},$ $-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$
98	$[-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},$ $\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{k},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},$ $-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},$ $\boldsymbol{j},\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{i},-\boldsymbol{j}]$

The **q** sequences have just a single $\boldsymbol{k}$, so for long lengths the frequency of each member of the alphabet asymptotically approaches a sequence over 4 roots of unity. With detailed knowledge of this approach to constructing long perfect quaternion sequences, Acevedo and Hall extended this result by noticing a connection to Lee sequences[9]. Acevedo and Hall chose not to reference this work[1].

2 Quaternion sequences with the AOP

Over roots of unity, a number of perfect sequence constructions exist which possess the AOP. These include the constructions of Heimiller-Frank[6][5], Milewski[11], the generalised sequence construction of Mow[13], and a construction by the author[4]. The existence of perfect quaternion sequences with the AOP has not been previously considered.

We considered a construction of the form

$$\mathbf{S} = [S_{a,b}] = \boldsymbol{i}^{\lfloor \frac{f(a,b)}{c} \rfloor} \boldsymbol{j}^{\lfloor \frac{g(a,b)}{d} \rfloor},$$

where $f(a,b)$ and $g(a,b)$ are bivariate polynomials with integer coefficients and c, d are positive integers. The sequence is formed by enumerating row-by-row the array $\mathbf{S}$. The size of the arrays considered were $2 \leq a, b \leq 32$, such that $ab > 16$. The integer coefficients of the polynomials and the denominators were less than 13. For each array size, 10 000 randomly selected polynomials and denominators were checked for the AOP.

The results of the search were interesting - one sequence of length 64 was found with the AOP. It was constructed from an array of size 8×8 with the index function

$$\mathbf{S} = [S_{a,b}] = \boldsymbol{i}^{a\,b} \boldsymbol{j}^{\lfloor \frac{a\,b}{2} \rfloor}.$$

The array and sequence is given by

$$\begin{bmatrix}
1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\
1 & \boldsymbol{i} & -\boldsymbol{j} & -\boldsymbol{k} & -1 & -\boldsymbol{i} & \boldsymbol{j} & \boldsymbol{k} \\
1 & -\boldsymbol{j} & -1 & \boldsymbol{j} & 1 & -\boldsymbol{j} & -1 & \boldsymbol{j} \\
1 & -\boldsymbol{k} & \boldsymbol{j} & \boldsymbol{i} & -1 & \boldsymbol{k} & -\boldsymbol{j} & -\boldsymbol{i} \\
1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\
1 & -\boldsymbol{i} & -\boldsymbol{j} & \boldsymbol{k} & -1 & \boldsymbol{i} & \boldsymbol{j} & -\boldsymbol{k} \\
1 & \boldsymbol{j} & -1 & -\boldsymbol{j} & 1 & \boldsymbol{j} & -1 & -\boldsymbol{j} \\
1 & \boldsymbol{k} & \boldsymbol{j} & -\boldsymbol{i} & -1 & -\boldsymbol{k} & -\boldsymbol{j} & \boldsymbol{i}
\end{bmatrix}$$

and

$$[1,1,1,1,1,1,1,1,1,\boldsymbol{i},-\boldsymbol{j},-\boldsymbol{k},-1,-\boldsymbol{i},\boldsymbol{j},\boldsymbol{k},1,-\boldsymbol{j},-1,\boldsymbol{j},1,-\boldsymbol{j},-1,$$
$$\boldsymbol{j},1,-\boldsymbol{k},\boldsymbol{j},\boldsymbol{i},-1,\boldsymbol{k},-\boldsymbol{j},-\boldsymbol{i},1,-1,1,-1,1,-1,1,-1,1,-\boldsymbol{i},$$
$$-\boldsymbol{j},\boldsymbol{k},-1,\boldsymbol{i},\boldsymbol{j},-\boldsymbol{k},1,\boldsymbol{j},-1,-\boldsymbol{j},1,\boldsymbol{j},-1,-\boldsymbol{j},1,\boldsymbol{k},\boldsymbol{j},-\boldsymbol{i},-1,-\boldsymbol{k},-\boldsymbol{j},\boldsymbol{i}].$$

The existence of this sequence is of interest, furthermore the fact that its length is the square of the number of distinct elements in the sequence draws parallels to the Heimiller-Frank construction. Based on the search not finding longer sequences with the AOP we make the following conjecture.

Conjecture 2.1. *The longest perfect sequence over the simple unit quaternions with the AOP is of length 64.*

3 Sequences of length 2^n, for $0 < n < 7$

CONSTRUCTION I Let $0 < n < 6$, we construct a sequence, $\mathbf{s} = [s_a]$, of length 2^n, where

$$s_a = \boldsymbol{i}^{\left\lfloor \frac{a^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2a^2}{2^{n-1}} \right\rfloor},$$

for $0 \leq a < 2^n$.

Theorem 3.1. *Let $\boldsymbol{s}$ be the sequence from Construction I. If τ is odd, then $\theta_{\boldsymbol{s}}(\tau) = 0$.*

Proof. The autocorrelation of $\mathbf{s}$ for shift τ is given by:

$$\begin{aligned}
\theta_{\mathbf{S}}(\tau) &= \sum_{a=0}^{2^n-1} s_a s_{a+\tau}^* \\
&= \sum_{a=0}^{2^n-1} \boldsymbol{i}^{\left\lfloor \frac{a^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2a^2}{2^{n-1}} \right\rfloor} \left(\boldsymbol{i}^{\left\lfloor \frac{(a+\tau)^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2(a+\tau)^2}{2^{n-1}} \right\rfloor} \right)^*
\end{aligned}$$

Over the quaternions, we have, $(ab)^* = b^* a^*$:

$$= \sum_{a=0}^{2^n-1} \boldsymbol{i}^{\left\lfloor \frac{a^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2a^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2(a+\tau)^2}{2^{n-1}} \right\rfloor^*} \boldsymbol{i}^{\left\lfloor \frac{(a+\tau)^2}{2^{n-1}} \right\rfloor^*}$$

Over the quaternions, we have, $q^* = |q|^2 q^{-1}$:

$$= \sum_{a=0}^{2^n-1} \boldsymbol{i}^{\left\lfloor \frac{a^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2a^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{-\left\lfloor \frac{2(a+\tau)^2}{2^{n-1}} \right\rfloor} \boldsymbol{i}^{-\left\lfloor \frac{(a+\tau)^2}{2^{n-1}} \right\rfloor}$$

Change coordinates by letting $a = q2^{n-1} + r$:

$$\begin{aligned}
&= \frac{1}{2} \sum_{q=0}^{3} \sum_{r=0}^{2^{n-1}-1} \boldsymbol{i}^{\left\lfloor \frac{q^2 2^{2n-2} + 2qr2^{n-1} + r^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2q^2 2^{2n-2} + 4qr2^{n-1} + 2r^2}{2^{n-1}} \right\rfloor} \times \\
&\boldsymbol{j}^{-\left\lfloor \frac{2q^2 2^{2n-2} + 4qr2^{n-1} + 2r^2 + 4\tau q2^{n-1} + 4\tau r + 2\tau^2}{2^{n-1}} \right\rfloor} \boldsymbol{i}^{-\left\lfloor \frac{q^2 2^{2n-2} + 2qr2^{n-1} + r^2 + 2\tau q2^{n-1} + 2\tau r + \tau^2}{2^{n-1}} \right\rfloor} \\
&= \frac{1}{2} \sum_{q=0}^{3} \sum_{r=0}^{2^{n-1}-1} \boldsymbol{i}^{2^{n-1}q^2 + 2qr + \left\lfloor \frac{r^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{2^n q^2 + 4qr + \left\lfloor \frac{2r^2}{2^{n-1}} \right\rfloor} \times \\
&\boldsymbol{j}^{-2^n q^2 - 4qr - 4\tau q - \left\lfloor \frac{2r^2 + 4\tau r + 2\tau^2}{2^{n-1}} \right\rfloor} \boldsymbol{i}^{-2^{n-1}q^2 - 2qr - 2\tau q - \left\lfloor \frac{r^2 + 2\tau r + \tau^2}{2^{n-1}} \right\rfloor} \\
&= \frac{1}{2} \sum_{q=0}^{3} \sum_{r=0}^{2^{n-1}-1} \boldsymbol{i}^{-2\tau q + \left\lfloor \frac{r^2}{2^{n-1}} \right\rfloor - \left\lfloor \frac{(r+\tau)^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{-4\tau q + \left\lfloor \frac{2r^2}{2^{n-1}} \right\rfloor - \left\lfloor \frac{2(r+\tau)^2}{2^{n-1}} \right\rfloor} \\
&= \frac{1}{2} \left(\sum_{q=0}^{3} \boldsymbol{i}^{-2\tau q} \right) \left(\sum_{r=0}^{2^{n-1}-1} \boldsymbol{i}^{\left\lfloor \frac{r^2}{2^{n-1}} \right\rfloor - \left\lfloor \frac{(r+\tau)^2}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{2r^2}{2^{n-1}} \right\rfloor - \left\lfloor \frac{2(r+\tau)^2}{2^{n-1}} \right\rfloor} \right) \qquad (1)
\end{aligned}$$

The leftmost summation in (1) is zero for τ odd. ∎

Remark 1. For $0 < n < 7$, we have confirmed by computer program that Construction I generates perfect sequences.

Example 3.2. We construct the 6 perfect sequences generated by Construction I. The sequences of length 2, 4, and 8 are Zadoff-Chu and Milewski sequences.

length 2: $[1, -\boldsymbol{i}]$
length 4: $[1, \boldsymbol{j}, -1, \boldsymbol{j}]$
length 8: $[1, 1, -\boldsymbol{i}, -1, 1, -1, -\boldsymbol{i}, 1]$
length 16: $[1, 1, \boldsymbol{j}, -\boldsymbol{i}, -1, \boldsymbol{i}, \boldsymbol{j}, -1, 1, -1, \boldsymbol{j}, \boldsymbol{i}, -1, -\boldsymbol{i}, \boldsymbol{j}, 1]$
length 32: $[1, 1, 1, \boldsymbol{j}, -\boldsymbol{i}, -\boldsymbol{k}, -1, \boldsymbol{i}, 1, -\boldsymbol{i}, -1, \boldsymbol{k}, -\boldsymbol{i}, -\boldsymbol{j}, 1,$
$-1, 1, -1, 1, -\boldsymbol{j}, -\boldsymbol{i}, \boldsymbol{k}, -1, -\boldsymbol{i}, 1, \boldsymbol{i}, -1, -\boldsymbol{k}, -\boldsymbol{i}, \boldsymbol{j}, 1, 1]$
length 64: $[1, 1, 1, 1, \boldsymbol{j}, \boldsymbol{j}, -\boldsymbol{i}, -\boldsymbol{k}, -1, -\boldsymbol{j}, \boldsymbol{i}, \boldsymbol{k}, \boldsymbol{j}, -\boldsymbol{i}, -1, \boldsymbol{i}, 1, -\boldsymbol{i}, -1, \boldsymbol{i}, \boldsymbol{j}, -\boldsymbol{k}, \boldsymbol{i}, \boldsymbol{j}, -1, \boldsymbol{k}, -\boldsymbol{i}, -\boldsymbol{j},$
$\boldsymbol{j}, -1, 1, -1, 1, -1, 1, -1, \boldsymbol{j}, -\boldsymbol{j}, -\boldsymbol{i}, \boldsymbol{k}, -1, \boldsymbol{j}, \boldsymbol{i}, -\boldsymbol{k}, \boldsymbol{j}, \boldsymbol{i}, -1, -\boldsymbol{i}, 1, \boldsymbol{i}, -1, -\boldsymbol{i}, \boldsymbol{j}, \boldsymbol{k}, \boldsymbol{i},$
$-\boldsymbol{j}, -1, -\boldsymbol{k}, -\boldsymbol{i}, \boldsymbol{j}, \boldsymbol{j}, 1, 1, 1]$

Extending Construction I to $n \geq 7$ generates good ZCZ sequences.

Example 3.3. Let $n = 7$, then Construction I generates the following sequence:
$[1, 1, 1, 1, 1, 1, \boldsymbol{j}, \boldsymbol{j}, -\boldsymbol{i}, -\boldsymbol{i}, -\boldsymbol{k}, -\boldsymbol{k}, -1, -\boldsymbol{j}, \boldsymbol{i}, \boldsymbol{k}, 1, \boldsymbol{j}, -\boldsymbol{i}, -\boldsymbol{k}, -1, -\boldsymbol{j}, \boldsymbol{k}, 1, -\boldsymbol{i}, -\boldsymbol{k}, -\boldsymbol{j}, \boldsymbol{i},$
$1, -\boldsymbol{i}, -1, \boldsymbol{i}, 1, -\boldsymbol{i}, -1, \boldsymbol{i}, 1, -\boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{k}, -\boldsymbol{i}, -1, \boldsymbol{k}, \boldsymbol{j}, -1, \boldsymbol{k}, -\boldsymbol{i}, -\boldsymbol{j}, 1, -\boldsymbol{k}, \boldsymbol{i}, \boldsymbol{j}, -1, \boldsymbol{k}, -\boldsymbol{k},$
$\boldsymbol{i}, -\boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{j}, -1, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, \boldsymbol{j}, -\boldsymbol{j}, -\boldsymbol{i}, \boldsymbol{i}, -\boldsymbol{k}, \boldsymbol{k}, -1, \boldsymbol{j}, \boldsymbol{i}, -\boldsymbol{k}, 1, -\boldsymbol{j}, -\boldsymbol{i}, \boldsymbol{k},$
$-1, \boldsymbol{j}, \boldsymbol{k}, -1, -\boldsymbol{i}, \boldsymbol{k}, -\boldsymbol{j}, -\boldsymbol{i}, 1, \boldsymbol{i}, -1, -\boldsymbol{i}, 1, \boldsymbol{i}, -1, -\boldsymbol{i}, 1, \boldsymbol{i}, -\boldsymbol{j}, -\boldsymbol{k}, -\boldsymbol{i}, 1, \boldsymbol{k}, -\boldsymbol{j}, -1, -\boldsymbol{k}, -\boldsymbol{i},$
$\boldsymbol{j}, 1, \boldsymbol{k}, \boldsymbol{i}, -\boldsymbol{j}, -1, -\boldsymbol{k}, -\boldsymbol{k}, -\boldsymbol{i}, -\boldsymbol{i}, \boldsymbol{j}, \boldsymbol{j}, 1, 1, 1, 1, 1],$

which has the following (left and right) autocorrelations:

128,0,0,0,0,0,0,0,16,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,16,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,-16,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,-16,0,0,0,0,0,0,0,0,0,0,0,0,0,0,-16,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,-16,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,16,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,16,0,0,0,0,0,0,0.

4 Arrays of size $2^n \times 2^n$, for $1 < n < 7$

CONSTRUCTION II Let $1 < n < 7$, we construct a 2-dimensional array, $\mathbf{S} = [S_{a,b}]$ of size $2^n \times 2^n$ over the unit quaternions: $\{-1, 1, -\boldsymbol{i}, \boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{j}, -\boldsymbol{k}, \boldsymbol{k}\}$, where

$$S_{a,b} = \boldsymbol{i}^{\left\lfloor \frac{4ab}{2^n} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{4a^2b^2}{2^n} \right\rfloor}.$$

Remark 2. For $1 < n < 7$, we have confirmed by computer program that Construction II generates perfect arrays.

Example 4.1. Let $n = 4$ as in Construction II, then we generate a 16×16 perfect array:

$$\left[\begin{array}{cccccccccccccccc}
1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\
1 & 1 & \boldsymbol{j} & -1 & \boldsymbol{i} & -\boldsymbol{i} & \boldsymbol{k} & \boldsymbol{i} & -1 & -1 & -\boldsymbol{j} & 1 & -\boldsymbol{i} & \boldsymbol{i} & -\boldsymbol{k} & -\boldsymbol{i} \\
1 & \boldsymbol{j} & \boldsymbol{i} & \boldsymbol{k} & -1 & -\boldsymbol{j} & -\boldsymbol{i} & -\boldsymbol{k} & 1 & \boldsymbol{j} & \boldsymbol{i} & \boldsymbol{k} & -1 & -\boldsymbol{j} & -\boldsymbol{i} & -\boldsymbol{k} \\
1 & -1 & \boldsymbol{k} & -1 & -\boldsymbol{i} & -\boldsymbol{i} & \boldsymbol{j} & -\boldsymbol{i} & -1 & 1 & -\boldsymbol{k} & 1 & \boldsymbol{i} & \boldsymbol{i} & -\boldsymbol{j} & \boldsymbol{i} \\
1 & \boldsymbol{i} & -1 & -\boldsymbol{i} & 1 & \boldsymbol{i} & -1 & -\boldsymbol{i} & 1 & \boldsymbol{i} & -1 & -\boldsymbol{i} & 1 & \boldsymbol{i} & -1 & -\boldsymbol{i} \\
1 & -\boldsymbol{i} & -\boldsymbol{j} & -\boldsymbol{i} & \boldsymbol{i} & -1 & -\boldsymbol{k} & -1 & -1 & \boldsymbol{i} & \boldsymbol{j} & \boldsymbol{i} & -\boldsymbol{i} & 1 & \boldsymbol{k} & 1 \\
1 & \boldsymbol{k} & -\boldsymbol{i} & \boldsymbol{j} & -1 & -\boldsymbol{k} & \boldsymbol{i} & -\boldsymbol{j} & 1 & \boldsymbol{k} & -\boldsymbol{i} & \boldsymbol{j} & -1 & -\boldsymbol{k} & \boldsymbol{i} & -\boldsymbol{j} \\
1 & \boldsymbol{i} & -\boldsymbol{k} & -\boldsymbol{i} & -\boldsymbol{i} & -1 & -\boldsymbol{j} & 1 & -1 & -\boldsymbol{i} & \boldsymbol{k} & \boldsymbol{i} & \boldsymbol{i} & 1 & \boldsymbol{j} & -1 \\
1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\
1 & -1 & \boldsymbol{j} & 1 & \boldsymbol{i} & \boldsymbol{i} & \boldsymbol{k} & -\boldsymbol{i} & -1 & 1 & -\boldsymbol{j} & -1 & -\boldsymbol{i} & -\boldsymbol{i} & -\boldsymbol{k} & \boldsymbol{i} \\
1 & -\boldsymbol{j} & \boldsymbol{i} & -\boldsymbol{k} & -1 & \boldsymbol{j} & -\boldsymbol{i} & \boldsymbol{k} & 1 & -\boldsymbol{j} & \boldsymbol{i} & -\boldsymbol{k} & -1 & \boldsymbol{j} & -\boldsymbol{i} & \boldsymbol{k} \\
1 & 1 & \boldsymbol{k} & 1 & -\boldsymbol{i} & \boldsymbol{i} & \boldsymbol{j} & \boldsymbol{i} & -1 & -1 & -\boldsymbol{k} & -1 & \boldsymbol{i} & -\boldsymbol{i} & -\boldsymbol{j} & -\boldsymbol{i} \\
1 & -\boldsymbol{i} & -1 & \boldsymbol{i} & 1 & -\boldsymbol{i} & -1 & \boldsymbol{i} & 1 & -\boldsymbol{i} & -1 & \boldsymbol{i} & 1 & -\boldsymbol{i} & -1 & \boldsymbol{i} \\
1 & \boldsymbol{i} & -\boldsymbol{j} & \boldsymbol{i} & \boldsymbol{i} & 1 & -\boldsymbol{k} & 1 & -1 & -\boldsymbol{i} & \boldsymbol{j} & -\boldsymbol{i} & -\boldsymbol{i} & -1 & \boldsymbol{k} & -1 \\
1 & -\boldsymbol{k} & -\boldsymbol{i} & -\boldsymbol{j} & -1 & \boldsymbol{k} & \boldsymbol{i} & \boldsymbol{j} & 1 & -\boldsymbol{k} & -\boldsymbol{i} & -\boldsymbol{j} & -1 & \boldsymbol{k} & \boldsymbol{i} & \boldsymbol{j} \\
1 & -\boldsymbol{i} & -\boldsymbol{k} & \boldsymbol{i} & -\boldsymbol{i} & 1 & -\boldsymbol{j} & -1 & -1 & \boldsymbol{i} & \boldsymbol{k} & -\boldsymbol{i} & \boldsymbol{i} & -1 & \boldsymbol{j} & 1
\end{array}\right]$$

5 Arrays of size $2^{n+1} \times 2^{n+1} \times 2^{n+1} \times 2^{n+1}$, for $0 < n < 6$

We now state a construction for 4-dimensional arrays over the unit quaternions.

CONSTRUCTION III Let $0 < n < 6$, we construct a 4-dimensional array, $\mathbf{S} = [S_{a,b,c,d}]$ of size $2^{n+1} \times 2^{n+1} \times 2^{n+1} \times 2^{n+1}$ over the unit quaternions: $\{-1, 1, -\boldsymbol{i}, \boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{j}, -\boldsymbol{k}, \boldsymbol{k}\}$, where

$$S_{a,b,c,d} = \boldsymbol{i}^{\left\lfloor \frac{ab}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{bc}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{cd}{2^{n-1}} \right\rfloor},$$

for $0 \leq a, b, c, d < 2^{n+1}$.

Theorem 5.1. *Let $\boldsymbol{S}$ be the array from Construction III. If $s_0 \neq 0 \mod 4$ or $s_3 \neq 0 \mod 4$, then $\theta_{\boldsymbol{S}}(s_0, s_1, s_2, s_3) = 0$.*

Proof. The autocorrelation of $\mathbf{S}$ for shift s_0, s_1, s_2, s_3 is given by:

$$\begin{aligned}
&\theta_{\mathbf{S}}(s_0, s_1, s_2, s_3) \\
&= \sum_{a=0}^{2^{n+1}-1} \sum_{b=0}^{2^{n+1}-1} \sum_{c=0}^{2^{n+1}-1} \sum_{d=0}^{2^{n+1}-1} S_{a,b,c,d}\, S^*_{a+s_0,b+s_1,c+s_2,d+s_3} \\
&= \sum_{a=0}^{2^{n+1}-1} \sum_{b=0}^{2^{n+1}-1} \sum_{c=0}^{2^{n+1}-1} \sum_{d=0}^{2^{n+1}-1} \boldsymbol{i}^{\left\lfloor \frac{ab}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{bc}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{cd}{2^{n-1}} \right\rfloor} \left(\boldsymbol{i}^{\left\lfloor \frac{(a+s_0)(b+s_1)}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{(b+s_1)(c+s_2)}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{(c+s_2)(d+s_3)}{2^{n-1}} \right\rfloor} \right)^*
\end{aligned}$$

Over the quaternions, we have, $(ab)^* = b^*a^*$:

$$= \sum_{a=0}^{2^{n+1}-1} \sum_{b=0}^{2^{n+1}-1} \sum_{c=0}^{2^{n+1}-1} \sum_{d=0}^{2^{n+1}-1} \boldsymbol{i}^{\left\lfloor \frac{ab}{2^{n-1}} \right\rfloor} \boldsymbol{j}^{\left\lfloor \frac{bc}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{cd}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{(c+s_2)(d+s_3)}{2^{n-1}} \right\rfloor^*} \boldsymbol{j}^{\left\lfloor \frac{(b+s_1)(c+s_2)}{2^{n-1}} \right\rfloor^*} \boldsymbol{i}^{\left\lfloor \frac{(a+s_0)(b+s_1)}{2^{n-1}} \right\rfloor^*} \tag{2}$$

Change coordinates by letting $a = q_a 2^{n-1} + r_a$, $b = q_b 2^{n-1} + r_b$, $c = q_c 2^{n-1} + r_c$, and $d = q_d 2^{n-1} + r_d$. Then (2) becomes:

$$\theta_{\mathbf{S}}(s_0, s_1, s_2, s_3) = \sum_{q_a=0}^{3} \sum_{r_a=0}^{2^{n-1}-1} \sum_{q_b=0}^{3} \sum_{r_b=0}^{2^{n-1}-1} \sum_{q_c=0}^{3} \sum_{r_c=0}^{2^{n-1}-1} \sum_{q_d=0}^{3} \sum_{r_d=0}^{2^{n-1}-1} \boldsymbol{i}^{\left\lfloor \frac{(q_a 2^{n-1}+r_a)(q_b 2^{n-1}+r_b)}{2^{n-1}} \right\rfloor} \times$$

$$\boldsymbol{j}^{\left\lfloor \frac{(q_b 2^{n-1}+r_b)(q_c 2^{n-1}+r_c)}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{(q_c 2^{n-1}+r_c)(q_d 2^{n-1}+r_d)}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{(q_c 2^{n-1}+r_c+s_2)(q_d 2^{n-1}+r_d+s_3)}{2^{n-1}} \right\rfloor^*} \times$$

$$\boldsymbol{j}^{\left\lfloor \frac{(q_b 2^{n-1}+r_b+s_1)(q_c 2^{n-1}+r_c+s_2)}{2^{n-1}} \right\rfloor^*} \boldsymbol{i}^{\left\lfloor \frac{(q_a 2^{n-1}+r_a+s_0)(q_b 2^{n-1}+r_b+s_1)}{2^{n-1}} \right\rfloor^*}$$

Expanding out the products, simplifying the *floor* functions, and factoring sums of products to products of sums yields:

$$\theta_{\mathbf{S}}(s_0, s_1, s_2, s_3) = \left(\sum_{q_a=0}^{3} \boldsymbol{i}^{-s_1 q_a} \right) \left(\sum_{q_b=0}^{3} \boldsymbol{i}^{-s_0 q_b} \boldsymbol{j}^{-s_2 q_b} \right) \left(\sum_{q_c=0}^{3} \boldsymbol{j}^{-s_1 q_c} \boldsymbol{k}^{-s_3 q_c} \right) \left(\sum_{q_d=0}^{3} \boldsymbol{k}^{-s_2 q_d} \right) \times$$

$$\left(\sum_{r_a=0}^{2^{n-1}-1} \sum_{r_b=0}^{2^{n-1}-1} \sum_{r_c=0}^{2^{n-1}-1} \sum_{r_d=0}^{2^{n-1}-1} \boldsymbol{i}^{\left\lfloor \frac{r_a r_b}{2^{n-1}} \right\rfloor - \left\lfloor \frac{r_a r_b + s_0 r_b + s_1 r_a + s_0 s_1}{2^{n-1}} \right\rfloor} \times \right.$$

$$\left. \boldsymbol{j}^{\left\lfloor \frac{r_b r_c}{2^{n-1}} \right\rfloor - \left\lfloor \frac{r_b r_c + s_1 r_c + s_2 r_b + s_1 s_2}{2^{n-1}} \right\rfloor} \boldsymbol{k}^{\left\lfloor \frac{r_c r_d}{2^{n-1}} \right\rfloor - \left\lfloor \frac{r_c r_d + s_2 r_d + s_3 r_c + s_2 s_3}{2^{n-1}} \right\rfloor} \right) \tag{3}$$

By Lemma 0.2, the first summation in (3) is zero for $s_1 \neq 0 \bmod 4$, similarly the fourth summation in (3) is zero for $s_2 \neq 0 \bmod 4$. Otherwise, for $s_1, s_2 = 0 \bmod 4$, the second and third summations in (3) are zero for $s_0, s_3 \neq 0 \bmod 4$. ∎

Remark 3. For $0 < n < 6$, Construction III generates perfect arrays.

We have confirmed by a computer program that the fifth summation in (3) is zero for $0 < n < 6$ and for all shifts, and non-zero for many shifts for $n = 6$.

Example 5.2. We generate the smallest perfect array from Construction III of size $4 \times 4 \times$

4×4:

$$\left[\begin{array}{cccc}
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & \boldsymbol{k} & -1 & -\boldsymbol{k} \\ 1 & -1 & 1 & -1 \\ 1 & -\boldsymbol{k} & -1 & \boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ \boldsymbol{j} & \boldsymbol{i} & -\boldsymbol{j} & -\boldsymbol{i} \\ -1 & 1 & -1 & 1 \\ -\boldsymbol{j} & \boldsymbol{i} & \boldsymbol{j} & -\boldsymbol{i} \end{array}\right] &
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ -1 & -\boldsymbol{k} & 1 & \boldsymbol{k} \\ 1 & -1 & 1 & -1 \\ -1 & \boldsymbol{k} & 1 & -\boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ -\boldsymbol{j} & -\boldsymbol{i} & \boldsymbol{j} & \boldsymbol{i} \\ -1 & 1 & -1 & 1 \\ \boldsymbol{j} & -\boldsymbol{i} & -\boldsymbol{j} & \boldsymbol{i} \end{array}\right] \\
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & \boldsymbol{k} & -1 & -\boldsymbol{k} \\ 1 & -1 & 1 & -1 \\ 1 & -\boldsymbol{k} & -1 & \boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} \boldsymbol{i} & \boldsymbol{i} & \boldsymbol{i} & \boldsymbol{i} \\ \boldsymbol{k} & -1 & -\boldsymbol{k} & 1 \\ -\boldsymbol{i} & \boldsymbol{i} & -\boldsymbol{i} & \boldsymbol{i} \\ -\boldsymbol{k} & -1 & \boldsymbol{k} & 1 \end{array}\right] &
\left[\begin{array}{cccc} -1 & -1 & -1 & -1 \\ 1 & \boldsymbol{k} & -1 & -\boldsymbol{k} \\ -1 & 1 & -1 & 1 \\ 1 & -\boldsymbol{k} & -1 & \boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} -\boldsymbol{i} & -\boldsymbol{i} & -\boldsymbol{i} & -\boldsymbol{i} \\ \boldsymbol{k} & -1 & -\boldsymbol{k} & 1 \\ \boldsymbol{i} & -\boldsymbol{i} & \boldsymbol{i} & -\boldsymbol{i} \\ -\boldsymbol{k} & -1 & \boldsymbol{k} & 1 \end{array}\right] \\
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & \boldsymbol{k} & -1 & -\boldsymbol{k} \\ 1 & -1 & 1 & -1 \\ 1 & -\boldsymbol{k} & -1 & \boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} -1 & -1 & -1 & -1 \\ -\boldsymbol{j} & -\boldsymbol{i} & \boldsymbol{j} & \boldsymbol{i} \\ 1 & -1 & 1 & -1 \\ \boldsymbol{j} & -\boldsymbol{i} & -\boldsymbol{j} & \boldsymbol{i} \end{array}\right] &
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ -1 & -\boldsymbol{k} & 1 & \boldsymbol{k} \\ 1 & -1 & 1 & -1 \\ -1 & \boldsymbol{k} & 1 & -\boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} -1 & -1 & -1 & -1 \\ \boldsymbol{j} & \boldsymbol{i} & -\boldsymbol{j} & -\boldsymbol{i} \\ 1 & -1 & 1 & -1 \\ -\boldsymbol{j} & \boldsymbol{i} & \boldsymbol{j} & -\boldsymbol{i} \end{array}\right] \\
\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & \boldsymbol{k} & -1 & -\boldsymbol{k} \\ 1 & -1 & 1 & -1 \\ 1 & -\boldsymbol{k} & -1 & \boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} -\boldsymbol{i} & -\boldsymbol{i} & -\boldsymbol{i} & -\boldsymbol{i} \\ -\boldsymbol{k} & 1 & \boldsymbol{k} & -1 \\ \boldsymbol{i} & -\boldsymbol{i} & \boldsymbol{i} & -\boldsymbol{i} \\ \boldsymbol{k} & 1 & -\boldsymbol{k} & -1 \end{array}\right] &
\left[\begin{array}{cccc} -1 & -1 & -1 & -1 \\ 1 & \boldsymbol{k} & -1 & -\boldsymbol{k} \\ -1 & 1 & -1 & 1 \\ 1 & -\boldsymbol{k} & -1 & \boldsymbol{k} \end{array}\right] &
\left[\begin{array}{cccc} \boldsymbol{i} & \boldsymbol{i} & \boldsymbol{i} & \boldsymbol{i} \\ -\boldsymbol{k} & 1 & \boldsymbol{k} & -1 \\ -\boldsymbol{i} & \boldsymbol{i} & -\boldsymbol{i} & \boldsymbol{i} \\ \boldsymbol{k} & 1 & -\boldsymbol{k} & -1 \end{array}\right]
\end{array}\right]$$

6 Arrays of size $2^n \times 2^n \times 2^{n+1} \times 2^{n+1}$, for $0 < n < 6$

We now state a second construction for 4-dimensional arrays over the unit quaternions. The construction is very similar to Construction III.

CONSTRUCTION IV Let $n > 0$ such that $n = 0 \bmod 4$, we construct a 4-dimensional array, $\mathbf{S} = [S_{a,b,c,d}]$ of size $2^n \times 2^n \times 2^{n+1} \times 2^{n+1}$ over the unit quaternions: $\{-1, 1, -\boldsymbol{i}, \boldsymbol{i}, -\boldsymbol{j}, \boldsymbol{j}, -\boldsymbol{k}, \boldsymbol{k}\}$, where

$$S_{a,b,c,d} = \boldsymbol{i}^{\lfloor \frac{ab}{2^{n-1}} \rfloor} \boldsymbol{j}^{\lfloor \frac{bc}{2^{n-1}} \rfloor} \boldsymbol{k}^{\lfloor \frac{cd}{2^{n-1}} \rfloor},$$

for $0 \leq a, b, c, d < 2^{n+1}$.

Remark 4. For $0 < n < 6$, we have confirmed by a computer program that Construction IV generates perfect arrays.

In this paper we have found, by exhaustive computer search, perfect sequences over the simple unit quaternions of lengths 10, 14, 18, 26, 30, 38, 42, 50, 54, 62, 74, 82, 90, and 98. These sequences were subsequently generalised to unbounded lengths via the Lee sequences. We have shown that perfect quaternion sequences with the AOP exist, and subsequently perfect sequences and arrays exist over the simple unit quaternions. We have conjectured that the Frank bound of the square of the number of elements in the alphabet of a perfect sequence extends to simple unit quaternions (Conjecture 2.1). We have discovered constructions for perfect 2 and 4-dimensional arrays over the simple unit quaternions (Construction I, Construction III, and Construction IV).

References

[1] S. B. Acevedo, T. E. Hall, "Perfect Sequences of Unbounded Lengths over the basic quaternions", *SETA 2012*, pp. 159-167, 2012

[2] S. B. Acevedo, N. Jolly, "Perfect arrays of unbounded size over the basic quaternions", *Cryptography and Communications*, vol. 6, no. 1, pp. 47-57, 2013

[3] K. T. Arasu, W. de Launey, "Two-dimensional perfect quaternary arrays", *IEEE Trans. Inf. Theory*, vol. 47, no. 4, pp. 1482-1493, 2001

[4] S. Blake, A. Z. Tirkel, "A Construction for Perfect Autocorrelation Sequences over Roots of Unity", *SETA, 2014*, pp. 104-108, November 2014

[5] R. L. Frank, S. A. Zadoff and R. Heimiller, "Phase Shift Pulse Codes with Good Periodic Correlation Properties", *IRE Transactions on Information Theory*, vol. 8, no. 6, pp. 381-382, October 1961

[6] R. C. Heimiller, "Phase Shift Pulse Codes with Good Periodic Correlation Properties", *IRE Transactions on Information Theory*, vol. 7, no. 4, pp. 254-257, October 1961

[7] O. Kuznetsov, "Perfect Sequences over the Real Quaternions", PhD thesis, School of Mathematical Sciences, Monash University, 2010

[8] O. Kuznetsov, T. Hall, "Perfect sequences over the real quaternions of longer length", *The Online Journal on Mathematics and Statistics*, pp. 17-20, Egypt, July 2010

[9] C. E. Lee, "Perfect q-ary sequences from multiplicative characters over GF(p)", *Electronic Letters 28*, pp. 833-835, 1992

[10] A. N. Leukhin, I. I. Popov, "Information compression by using polarization properties of photon echo", *IRQO '99: Quantum Optics, Vitali V. Samartsev; Ed.*, Proc. SPIE, vol. 4061, p. 85-91

[11] A. Milewski, "Periodic Sequences with Optimal Properties for Channel Estimation and Fast Start-Up Equalization", *IBM Journal of Research and Development*, vol. 27, no. 5, pp. 426-431, September 1983

[12] Monash Campus Grid, Monash University, Clayton, Australia. http://www.monash.edu.au/eresearch/services/mcg/

[13] W. H. Mow, "A New Unified Construction of Perfect Root-of-Unity Sequences", *IEEE 4th Inter. Symp. on Spread Spectrum Techniques and Applications Proceedings*, vol. 3, pp. 955-959, 1996

[14] C. E. Moxey, T. A. Ell, S. J. Sangwine, "Hypercomplex operators and vector correlation.", *Proc. XI Eur. Signal Process. Conf. Toulouse, France*, 2002

[15] S. Pei, J. Ding, J. Chang, "Efficient implementation of quaternion Fourier transform, convolution, and correlation by 2-D complex FFT", *IEEE trans. on Signal Processing*, pp. 2783-2797, November 2001

[16] S. J. Sangwine, T. A. Ell, "Hypercomplex Auto- and Cross-Correlation of Color Images", *Poster presented at IEEE International Conference on Image Processing (ICIP99)*, Kobe, Japan, Thursday 28 October 1999

[17] K. Smith-Miles, letter from Monash university, 2013